

Image










LRIG3 Antibody

Product Code	CSB-PA751017LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q6UXM1
Immunogen	Recombinant Human Leucine-rich repeats and immunoglobulin-like domains protein 3 protein (835-985AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IF; Recommended dilution: IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
	Dragger (office) 0.000/ Dragglin 200
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Storage Buffer Purification Method	
	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified
Purification Method Isotype	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG
Purification Method Isotype Clonality	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG Polyclonal Leucine-rich repeats and immunoglobulin-like domains protein 3, LIG-3, LRIG3,
Purification Method Isotype Clonality Alias	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG Polyclonal Leucine-rich repeats and immunoglobulin-like domains protein 3, LIG-3, LRIG3, LIG3, UNQ287/PRO326/PRO335
Purification Method Isotype Clonality Alias Immunogen Species	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG Polyclonal Leucine-rich repeats and immunoglobulin-like domains protein 3, LIG-3, LRIG3, LIG3, UNQ287/PRO326/PRO335 Homo sapiens (Human)



Immunofluorescence staining of U251 cells with CSB-PA751017LA01HU at 1:100, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).