




DNAJA3 Antibody

Product Code	CSB-PA836207LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q96EY1
Immunogen	Recombinant Human DnaJ homolog subfamily A member 3, mitochondrial protein (86-184AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IF, IP; Recommended dilution: WB:1:1000-1:5000, IHC:1:200-1:500, IF:1:50-1:200, IP:1:200-1:2000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	DnaJ homolog subfamily A member 3, mitochondrial (DnaJ protein Tid-1) (hTid-1) (Hepatocellular carcinoma-associated antigen 57) (Tumorous imaginal discs protein Tid56 homolog), DNAJA3, HCA57 TID1
Immunogen Species	Homo sapiens (Human)
Research Area	Signal Transduction
Target Names	DNAJA3

Image




Western Blot


Positive WB detected in: HeLa whole cell lysate, MCF-7 whole cell lysate, K562 whole cell lysate, Jurkat whole cell lysate, Raji whole cell lysate, 293T whole cell lysate, A549 whole cell lysate
All lanes: DNAJA3 antibody at 1:2000

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution
Predicted band size: 53, 50, 34 kDa
Observed band size: 53 kDa




IHC image of CSB-PA836207LA01HU diluted at 1:400 and staining in paraffin-embedded human lung cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA836207LA01HU diluted at 1:400 and staining in paraffin-embedded human pancreatic cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of MCF-7 cells with CSB-PA836207LA01HU at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunoprecipitating DNAJA3 in HeLa whole cell lysate
 Lane 1: Rabbit control IgG instead of CSB-PA836207LA01HU in HeLa whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/50000)
 Lane 2: CSB-PA836207LA01HU (5μg) + HeLa whole cell lysate (0.5mg)
 Lane 3: HeLa whole cell lysate (20μg)