




# TMEM214 Antibody

<b>Product Code</b>	CSB-PA023808LA01HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q6NUQ4
<b>Immunogen</b>	Recombinant Human Transmembrane protein 214 protein (2-236AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IP; Recommended dilution: WB:1:1000-1:5000, IP:1:200-1:2000
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
<b>Purification Method</b>	>95%, Protein G purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Transmembrane protein 214, TMEM214
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Cell Biology
<b>Target Names</b>	TMEM214

## Image



### Western Blot

Positive WB detected in: K562 whole cell lysate, HepG2 whole cell lysate, HL-60 whole cell lysate, A549 whole cell lysate, U251 whole cell lysate


All lanes: TMEM214 antibody at 1:2000

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 78, 73 kDa

Observed band size: 78 kDa



Immunoprecipitating TMEM214 in HepG2 whole cell lysate

Lane 1: Rabbit control IgG instead of CSB-PA023808LA01HU in HepG2 whole cell lysate.

For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/50000)

Lane 2: CSB-PA023808LA01HU (5ug) + HepG2 whole cell lysate (0.5mg)

Lane 3: HepG2 whole cell lysate (20ug)