




KAT2A Antibody

| | |
|----------------------------|---|
| Product Code | CSB-PA821706LA01HU |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q92830 |
| Immunogen | Recombinant Human Histone acetyltransferase KAT2A protein (302-537AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, IHC; Recommended dilution: IHC:1:200-1:500 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Histone acetyltransferase KAT2A (EC 2.3.1.48) (General control of amino acid synthesis protein 5-like 2) (Histone acetyltransferase GCN5) (hGCN5) (Histone succinyltransferase KAT2A) (EC 2.3.1.-) (Lysine acetyltransferase 2A) (STAF97), KAT2A, GCN5 GCN5L2 |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Epigenetics and Nuclear Signaling |
| Target Names | KAT2A |

Image



IHC image of CSB-PA821706LA01HU diluted at 1:300 and staining in paraffin-embedded human melanoma cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.