




CD14 Monoclonal Antibody

Product Code	CSB-MA004879A1m
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P08571
Immunogen	Recombinant Human Monocyte differentiation antigen CD14 protein (20-345AA)
Raised In	Mouse
Species Reactivity	Human, Mouse, Rat, Rabbit
Tested Applications	ELISA, WB, IF; Recommended dilution: WB:1:1000-1:64000, IF:1:50-1:200
Relevance	Coreceptor for bacterial lipopolysaccharide (PubMed:1698311, PubMed:23264655). In concert with LBP, binds to monomeric lipopolysaccharide and delivers it to the LY96/TLR4 complex, thereby mediating the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed:20133493, PubMed:23264655). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:8612135). Acts as a coreceptor for TLR2:TLR6 heterodimer in response to diacylated lipopeptides and for TLR2:TLR1 heterodimer in response to triacylated lipopeptides, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway (PubMed:16880211). Binds electronegative LDL (LDL-) and mediates the cytokine release induced by LDL- (PubMed:23880187).
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein A purified
Isotype	IgG1
Clonality	Monoclonal
Product Type	Monoclonal Antibody
Immunogen Species	Homo sapiens (Human)
Target Names	CD14
Clone No.	1A7C5
Image	



Western Blot

Positive WB detected in: Rabbit lung tissue, Rabbit liver tissue, Rabbit spleen tissue, Rabbit small intestine tissue


All lanes: CD14 antibody at 1:2500

Secondary

Goat polyclonal to Mouse IgG at 1/10000 dilution

Predicted band size: 41 kDa

Observed band size: 55 kDa



Western Blot


Positive WB detected in: Mouse kidney tissue, Rat liver tissue

All lanes: CD14 antibody at 1:2000

Secondary Goat polyclonal to Mouse IgG at 1/10000 dilution

Predicted band size: 41 kDa

Observed band size: 55 kDa



Western Blot

Positive WB detected in: NIH/3T3 whole cell lysate


All lanes: CD14 antibody at at 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000, 1:64000

Secondary

Goat polyclonal to Mouse IgG at 1/10000 dilution

Predicted band size: 41 kDa

Observed band size: 55 kDa



Western Blot

Positive WB detected in: HepG2 whole cell lysate, A549 whole cell lysate, HeLa whole cell lysate, Jurkat whole cell lysate, NIH/3T3 whole cell lysate


All lanes: CD14 antibody at 1:1000

Secondary


Goat polyclonal to Mouse IgG at 1/10000 dilution

Predicted band size: 41 kDa


Observed band size: 55 kDa



Immunofluorescence staining of A549 cells with CSB-MA004879A1m at 1:100, counter-stained with DAPI. The cells were blocked in 10% normal Goat Serum and then incubated with the primary antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescence staining of HepG2 cells with CSB-MA004879A1m at 1:100, counter-stained with DAPI. The cells were blocked in 10% normal Goat Serum and then incubated with the primary antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescence staining of Raw264.7 cells with CSB-MA004879A1m at 1:100, counter-stained with DAPI. The cells were blocked in 10% normal Goat Serum and then incubated with the primary antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.