


🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🌘

FZD7 Antibody

| Product Code | CSB-PA009110LA01HU |
|---------------------|--|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | O75084 |
| Immunogen | Recombinant Human Frizzled-7 protein (161-239AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:200-1:500 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method | >95%, Protein G purified |
| lsotype | lgG |
| Clonality | Polyclonal |
| Alias | Frizzled-7 (Fz-7) (hFz7) (FzE3), FZD7 |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Signal Transduction |
| Target Names | FZD7 |
| | |


Image



1



🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🕒 Website: www.cusabio.com 🧉



IHC image of CSB-PA009110LA01HU diluted at 1:400 and staining in paraffin-embedded human adrenal gland tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.