




GPR37 Antibody

Product Code	CSB-PA009810NA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	O15354
Immunogen	Peptide sequence from Human Prosaposin receptor GPR37 protein (114-133AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:1000-1:5000, IHC:1:500-1:1000, IF:1:200-1:500
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Prosaposin receptor GPR37 (Endothelin B receptor-like protein 1) (ETBR-LP-1) (G-protein coupled receptor 37) (Parkin-associated endothelin receptor-like receptor) (PAELR), GPR37
Immunogen Species	Homo sapiens (Human)
Research Area	Signal Transduction
Target Names	GPR37

Image




Western Blot


Positive WB detected in: 293 whole cell lysate, PC-3 whole cell lysate, MCF-7 whole cell lysate
All lanes: GPR37 antibody at 1:2000

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution
Predicted band size: 68 kDa
Observed band size: 68 kDa



IHC image of CSB-PA009810NA01HU diluted at 1:800 and staining in paraffin-embedded human liver tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of PC-3 cells with CSB-PA009810NA01HU at 1:266, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.