


🕜 Tel: +1-301-363-4651 🛛 🖾 Email: cusabio@cusabio.com 🛛 🥑 Website: www.cusabio.com 🌘


## **OPRK1** Antibody

Product Code	CSB-PA016359OA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P41145
Immunogen	Peptide sequence from Human Kappa-type opioid receptor protein (31-50AA)
Raised In	Rabbit
Species Reactivity	Human
<b>Tested Applications</b>	ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	Antigen Affinity Purified
Isotype	lgG
Clonality	Polyclonal
Alias	Kappa-type opioid receptor (K-OR-1) (KOR-1), OPRK1, OPRK
Immunogen Species	Homo sapiens (Human)
Research Area	Neuroscience
Target Names	OPRK1

Image



IHC image of CSB-PA016359OA01HU diluted at 1:100 and staining in paraffin-embedded human ovarian cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of HepG2 cells with CSB-PA016359OA01HU at 1:50, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).

1