




ARL2BP Antibody

| | |
|----------------------------|---|
| Product Code | CSB-PA897097HA01HU |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q9Y2Y0 |
| Immunogen | Recombinant Human ADP-ribosylation factor-like protein 2-binding protein (1-163AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method | Antigen Affinity Purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | ADP-ribosylation factor-like protein 2-binding protein (ARF-like 2-binding protein) (ARL2-binding protein) (Binder of ARF2 protein 1), ARL2BP, BART BART1 |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Signal Transduction |
| Target Names | ARL2BP |

Image



Western Blot

Positive WB detected in: HepG2 whole cell lysate, U251 whole cell lysate, U87 whole cell lysate


All lanes: ARL2BP antibody at 1:1000

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 19, 18 kDa

Observed band size: 19 kDa



IHC image of CSB-PA897097HA01HU diluted at 1:100 and staining in paraffin-embedded human glioma cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.