




# POGLUT1 Antibody

<b>Product Code</b>	CSB-PA818754LA01HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q8NBL1
<b>Immunogen</b>	Recombinant Human Protein O-glucosyltransferase 1 protein (132-392AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:200-1:500
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
<b>Purification Method</b>	>95%, Protein G purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Protein O-glucosyltransferase 1 (EC 2.4.1.-) (CAP10-like 46 kDa protein) (hCLP46) (KTEL motif-containing protein 1) (Myelodysplastic syndromes relative protein) (O-glucosyltransferase Rumi homolog) (hRumi) (Protein O-xylosyltransferase POGLUT1) (EC 2.4.2.26), POGLUT1, C3orf9 CLP46 KTELC1 MDSRP
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Developmental Biology
<b>Target Names</b>	POGLUT1

## Image



### Western Blot

Positive WB detected in: HepG2 whole cell lysate, PC-3 whole cell lysate, K562 whole cell lysate, 293 whole cell lysate


All lanes: POGLUT1 antibody at 1:2000

### Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 47 kDa

Observed band size: 47 kDa



IHC image of CSB-PA818754LA01HU diluted at 1:300 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

## Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.