




POGLUT1 Antibody

| | |
|----------------------------|--|
| Product Code | CSB-PA818754LA01HU |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q8NBL1 |
| Immunogen | Recombinant Human Protein O-glucosyltransferase 1 protein (132-392AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:200-1:500 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Protein O-glucosyltransferase 1 (EC 2.4.1.-) (CAP10-like 46 kDa protein) (hCLP46) (KTEL motif-containing protein 1) (Myelodysplastic syndromes relative protein) (O-glucosyltransferase Rumi homolog) (hRumi) (Protein O-xylosyltransferase POGLUT1) (EC 2.4.2.26), POGLUT1, C3orf9 CLP46 KTELC1 MDSRP |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Developmental Biology |
| Target Names | POGLUT1 |

Image



Western Blot

Positive WB detected in: HepG2 whole cell lysate, PC-3 whole cell lysate, K562 whole cell lysate, 293 whole cell lysate


All lanes: POGLUT1 antibody at 1:2000

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 47 kDa

Observed band size: 47 kDa



IHC image of CSB-PA818754LA01HU diluted at 1:300 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.