









## **ANKRD42** Antibody

| <b>Product Code</b>        | CSB-PA847661LA01HU  |
|----------------------------|---|
| Storage                    | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.                     |
| Uniprot No.                | Q8N9B4  |
| Immunogen                  | Recombinant Human Ankyrin repeat domain-containing protein 42 protein (158-377AA) |
| Raised In                  | Rabbit  |
| Species Reactivity         | Human   |
| <b>Tested Applications</b> | ELISA, IHC; Recommended dilution: IHC:1:500-1:5000                                |
| Form                       | Liquid  |
| Conjugate                  | Non-conjugated  |
| Storage Buffer             | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, pH 7.4  |
| Purification Method        | >95%, Antigen Affinity purified & Affinity purified                               |
| Isotype                    | IgG   |
| Clonality                  | Polyclonal  |
| Alias                      | Ankyrin repeat domain-containing protein 42, ANKRD42                              |
| Immunogen Species          | Homo sapiens (Human)  |
| Research Area              | Signal Transduction   |
| Target Names               | ANKRD42   |
| Image                      | ILIC image of CCD DA947CC41 A04LILI diluted at                                    |



IHC image of CSB-PA847661LA01HU diluted at 1:500 and staining in paraffin-embedded human prostate tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG labeled by HRP and visualized using 0.05% DAB.



IHC image of CSB-PA847661LA01HU diluted at 1:500 and staining in paraffin-embedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG labeled by HRP and visualized



## **CUSABIO TECHNOLOGY LLC**



using 0.05% DAB.