




# HSPA8 Monoclonal Antibody

|                            |   |
|----------------------------|---|
| <b>Product Code</b>        | CSB-MA010829A0m   |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |
| <b>Uniprot No.</b>         | P11142  |
| <b>Immunogen</b>           | Recombinant Human Heat shock cognate 71 kDa protein (2-646AA)   |
| <b>Raised In</b>           | Mouse   |
| <b>Species Reactivity</b>  | Human, Rat, Mouse   |
| <b>Tested Applications</b> | ELISA, WB, IHC, FC, IP; Recommended dilution: WB: 1:10000-1:256000, IHC: 1:100-1:500, FC: 1:100-1:300, IP: 1µl-4µl  |
| <b>Relevance</b>           | Acts as a repressor of transcriptional activation. Inhibits the transcriptional coactivator activity of CITED1 on Smad-mediated transcription. Chaperone. Component of the PRP19-CDC5L complex that forms an integral part of the spliceosome and is required for activating pre-mRNA splicing. May have a scaffolding role in the spliceosome assembly as it contacts all other components of the core complex. Binds bacterial lipopolysaccharide (LPS) et mediates LPS-induced inflammatory response, including TNF secretion by monocytes. Participates in the ER-associated degradation (ERAD) quality control pathway in conjunction with J domain-containing co-chaperones and the E3 ligase CHIP. |
| <b>Form</b>                | Liquid  |
| <b>Conjugate</b>           | Non-conjugated  |
| <b>Storage Buffer</b>      | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  |
| <b>Purification Method</b> | >95%, Protein A purified  |
| <b>Isotype</b>             | IgG2b   |
| <b>Clonality</b>           | Monoclonal  |
| <b>Alias</b>               | Heat shock cognate 71 kDa protein (Heat shock 70 kDa protein 8)<br>(Lipopolysaccharide-associated protein 1) (LAP-1) (LPS-associated protein 1),<br>HSPA8, HSC70 HSP73 HSPA10   |
| <b>Product Type</b>        | Monoclonal Antibody   |
| <b>Immunogen Species</b>   | Homo sapiens (Human)  |
| <b>Gene Names</b>          | HSPA8   |
| <b>Clone No.</b>           | 2G8F6   |
| <b>Image</b>               |   |



#### Western Blot

Positive WB detected in: HeLa whole cell lysate, NIH/3T3 whole cell lysate, K562 whole cell lysate, HepG2 whole cell lysate, Mouse spleen tissue, Rat spleen tissue, Mouse heart tissue, Rat heart tissue

All lanes HSPA8 antibody at 1:2000


#### Secondary

Goat polyclonal to mouse IgG at 1/50000 dilution

Predicted band size: 70~75 KDa

Observed band size: 70~75 KDa

Exposure time: 10s



#### Western Blot

Positive WB detected in: HeLa whole cell lysate at 20µg, 10µg, 5µg, 2.5µg, 1.25µg, 0.625µg

All lanes: HSPA8 antibody at 1:2000


#### Secondary

Goat polyclonal to mouse IgG at 1/50000 dilution

Predicted band size: 70~75 KDa

Observed band size: 70~75 KDa

Exposure time: 10s



#### Western Blot

Positive WB detected in: 20µg HeLa whole cell lysate

HSPA8 antibody at 1:2000, 1:4000, 1:8000, 1:16000, 1:32000, 1:64000, 1:128000, 1:256000


#### Secondary

Goat polyclonal to mouse IgG at 1/50000 dilution


Predicted band size: 70~75 KDa

Observed band size: 70~75 KDa

Exposure time: 10s




IHC image of CSB-MA010829A0m diluted at 1:256 and staining in paraffin-embedded human prostate cancer performed on a Leica Bond<sup>TM</sup> system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight, and detected by a Goat anti-mouse IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC image of CSB-MA010829A0m diluted at 1:256 and staining in paraffin-embedded human prostate cancer performed on a Leica Bond<sup>TM</sup> system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight, and detected by a Goat anti-mouse IgG polymer labeled by HRP and



visualized using 0.05% DAB.




Immunoprecipitating HSPA8 in HeLa whole cell lysate

Lane 1: Mouse control IgG2b instead of CSB-MA010829A0m in HeLa whole cell lysate

Lane 2: CSB-MA010829A0m (1.5μl) + HeLa whole cell lysate (500μg)

Lane 3: HeLa whole cell lysate (20μg)

For western blotting, the blot was detected with CSB-MA010829A0m at 1:2000, and a HRP-conjugated Protein G antibody was used as the secondary antibody at 1:2000



Overlay histogram showing MCF-7 cells stained with CSB-MA010829A0m (red line). The cells were fixed with 70% Ethylalcohol (18h) and then incubated in 10% normal goat serum to block non-specific protein-protein interactions followed by the primary antibody at 1:200 for 1 h at 4°C. The secondary antibody used was FITC goat anti-mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG2b used under the same conditions. Acquisition of >10,000 events was performed.