




# YWHAZ Monoclonal Antibody

<b>Product Code</b>	CSB-MA026293A0m
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	P63104
<b>Immunogen</b>	Recombinant Human 14-3-3 protein zeta/delta protein (133-212AA)
<b>Raised In</b>	Mouse
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Tested Applications</b>	ELISA, WB, IHC, IF, FC; Recommended dilution: WB:1:5000-160000, IHC:1:50-1:200, IF:1:50-1:200, FC:1:50-1:200, IP:1µl-4µl
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	>95%, Protein A purified
<b>Isotype</b>	IgG2b
<b>Clonality</b>	Monoclonal Antibody
<b>Product Type</b>	Monoclonal Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Target Names</b>	YWHAZ
<b>Clone No.</b>	19G7E9

## Image



### Western Blot

Positive WB detected in: NIH/3T3 whole cell lysate, HeLa whole cell lysate, HepG2 whole cell lysate, A549 whole cell lysate, RAW264.7 whole cell lysate, Rat Brain tissue

All lanes YWHAZ antibody at 1:5000


### Secondary

Goat polyclonal to mouse IgG at 1/50000 dilution

Predicted band size: 34 KDa

Observed band size: 34 KDa

Exposure time:1min



**Western Blot**

Positive WB detected in: HeLa whole cell lysate at 20µg, 10µg, 5µg, 2.5µg, 1.25µg, 0.625µg

All lanes: YWHAZ antibody at 1:5000


**Secondary**

Goat polyclonal to mouse IgG at 1/50000 dilution

Predicted band size: 34 KDa

Observed band size: 34 KDa

Exposure time:5min



**Western Blot**

Positive WB detected in: 20µg HeLa whole cell lysate YWHAZ antibody

at 1:5000, 1:10000, 1:20000, 1:40000, 1:80000, 1:160000


**Secondary**

Goat polyclonal to mouse IgG at 1/50000 dilution


Predicted band size: 34 KDa

Observed band size: 34 KDa


Exposure time:5min



IHC image of CSB-MA026293A0m diluted at 1:200 and staining in paraffin-embedded human lung cancer tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at 37°C. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG labeled by HRP and visualized using 0.05% DAB.




IHC image of CSB-MA026293A0m diluted at 1:200 and staining in paraffin-embedded human liver cancer tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at 37°C. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG labeled by HRP and visualized using 0.05% DAB.




IHC image of CSB-MA026293A0m diluted at 1:200 and staining in paraffin-embedded human breast cancer tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at 37°C. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG labeled by HRP and




visualized using 0.05% DAB.




Immunofluorescence staining of A549 cells with CSB-MA026293A0m at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).




Immunofluorescence staining of Hela cells with CSB-MA026293A0m at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).




Immunofluorescence staining of U251 cells with CSB-MA026293A0m at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).




Overlay histogram showing A549 cells stained with CSB-MA026293A0m (red line) at 1:100. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum was Incubated to block non-specific protein-protein interactions followed by the antibody (1µg/1\*10<sup>6</sup>cells) for 1 h at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG2b (1µg/1\*10<sup>6</sup>cells) used under the same conditions. Acquisition of >10,000 events was performed.




Overlay histogram showing HeLa cells stained with CSB-MA026293A0m (red line) at 1:100. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum was incubated to block non-specific protein-protein interactions followed by the antibody ( $1\mu\text{g}/1*10^6$  cells) for 1 h at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG2b ( $1\mu\text{g}/1*10^6$  cells) used under the same conditions. Acquisition of >10,000 events was performed.



Overlay histogram showing HepG2 cells stained with CSB-MA026293A0m (red line) at 1:100. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum was incubated to block non-specific protein-protein interactions followed by the antibody ( $1\mu\text{g}/1*10^6$  cells) for 1 h at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG2b ( $1\mu\text{g}/1*10^6$  cells) used under the same conditions. Acquisition of >10,000 events was performed.



Immunoprecipitating YWHAZ in A549 whole cell lysate  
 Lane 1: Mouse control IgG2b instead of CSB-MA026293A0m in A549 whole cell lysate.  
 Lane 2: CSB-MA026293A0m ( $1\mu\text{g}$ ) + A549 whole cell lysate ( $500\mu\text{g}$ )  
 Lane 3: A549 whole cell lysate ( $20\mu\text{g}$ )  
 For western blotting, the blot was detected with CSB-MA026293A0m at 1:2000, and a HRP-conjugated Protein G antibody was used as the secondary antibody at 1:50000



Immunoprecipitating YWHAZ in HepG2 whole cell lysate  
 Lane 1: Mouse control IgG2b instead of CSB-MA026293A0m in HepG2 whole cell lysate.  
 Lane 2: CSB-MA026293A0m ( $1\mu\text{g}$ ) + HepG2 whole cell lysate ( $500\mu\text{g}$ )  
 Lane 3: HepG2 whole cell lysate ( $20\mu\text{g}$ )  
 For western blotting, the blot was detected with CSB-MA026293A0m at 1:2000, and a HRP-conjugated Protein G antibody was used as the secondary antibody at 1:50000