




CD19 Recombinant Monoclonal Antibody

Product Code	CSB-RA780821A0HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P15391
Immunogen	A synthesized peptide derived from human CD19
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IP; Recommended dilution: WB:1:500-1:5000, IHC:1:50-1:200, IP:1:200-1:1000
Relevance	Assembles with the antigen receptor of B-lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Rabbit IgG in 10mM phosphate buffered saline , pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Purification Method	Affinity-chromatography
Isotype	Rabbit IgG
Clonality	Monoclonal
Product Type	Recombinant Antibody
Immunogen Species	Homo sapiens (Human)
Research Area	Immunology; Stem cells
Target Names	CD19
Clone No.	4C7

Image



Western Blot

Positive WB detected in: Daudi whole cell lysate, Raji whole cell lysate


All lanes: CD19 antibody at 1:2000

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 62 kDa

Observed band size: 95 kDa



IHC image of CSB-RA780821A0HU diluted at 1:100 and staining in paraffin-embedded human tonsil tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4? overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



Immunoprecipitating CD19 in Raji whole cell lysate

Lane 1: Rabbit control IgG instead of CSB-RA780821A0HU in Raji whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)

Lane 2: CSB-RA780821A0HU(2μg)+ Raji whole cell lysate(500μg)

Lane 3: Raji whole cell lysate (10μg)

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.