




CD63 Monoclonal Antibody

Product Code	CSB-MA004950A1m
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P08962
Immunogen	Recombinant Human CD63 antigen protein (103-203AA)
Raised In	Mouse
Species Reactivity	Human, Rabbit
Tested Applications	ELISA, WB, IHC, IF, FC; Recommended dilution: WB:1:1000-1:8000, IHC:1:50-1:200, IF:1:50-1:200, FC:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG1
Clonality	Monoclonal Antibody
Product Type	Monoclonal Antibody
Immunogen Species	Homo sapiens (Human)
Gene Names	CD63
Clone No.	10F11E6

Image



Western Blot

Positive WB detected in: A549 whole cell lysate, HeLa whole cell lysate, HepG2 whole cell lysate, MCF-7 whole cell lysate

All lanes CD63 antibody at 1:1000


Secondary

Goat polyclonal to mouse IgG at 1/50000 dilution

Predicted band size: 30-120 KD KDa

Observed band size: 30-120 KD KDa

Exposure time:1min



Western Blot

Positive WB detected in: Raji whole cell lysate

All lanes CD63 antibody at 1:1000


Secondary

Goat polyclonal to mouse IgG at 1/50000 dilution

Predicted band size: 30-120 KD KDa

Observed band size: 30-120 KD KDa

Exposure time:1min



Western Blot

Positive WB detected in: A375 whole cell lysate,

Rabbit spleen tissue

All lanes CD63 antibody at 1:1000


Secondary

Goat polyclonal to mouse IgG at 1/50000 dilution


Predicted band size: 30-120 KD KDa

Observed band size: 30-120 KD KDa


Exposure time:1min`




IHC image of CSB-MA004950A1m diluted at 1:500 and staining in paraffin-embedded human lung cancer tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at 37°C. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG labeled by HRP and visualized using 0.05% DAB.




IHC image of CSB-MA004950A1m diluted at 1:500 and staining in paraffin-embedded human lung cancer tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at 37°C. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG labeled by HRP and visualized using 0.05% DAB.




IHC image of CSB-MA004950A1m diluted at 1:500 and staining in paraffin-embedded human lung cancer tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at 37°C. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG labeled by HRP and visualized using 0.05% DAB.




Immunofluorescence staining of A549 cells with CSB-MA004950A1m at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).




Immunofluorescence staining of Hela cells with CSB-MA004950A1m at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).




Immunofluorescence staining of MCF-7 cells with CSB-MA004950A1m at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).




Overlay histogram showing A549 cells stained with CSB-MA004950A1m (red line) at 1:100. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum was incubated to block non-specific protein-protein interactions followed by the antibody (1µg/1*10⁶ cells) for 1 h at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG2b (1µg/1*10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.




Overlay histogram showing Hela cells stained with CSB-MA004950A1m (red line) at 1:100. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum was incubated to block non-specific protein-protein interactions followed by the antibody ($1\mu\text{g}/1 \times 10^6$ cells) for 1 h at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG2b ($1\mu\text{g}/1 \times 10^6$ cells) used under the same conditions. Acquisition of >10,000 events was performed.



Overlay histogram showing HepG2 cells stained with CSB-MA004950A1m (red line) at 1:100. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum was incubated to block non-specific protein-protein interactions followed by the antibody ($1\mu\text{g}/1 \times 10^6$ cells) for 1 h at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG2b ($1\mu\text{g}/1 \times 10^6$ cells) used under the same conditions. Acquisition of >10,000 events was performed.



Overlay histogram showing K562 cells stained with CSB-MA004950A1m (red line) at 1:100. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum was incubated to block non-specific protein-protein interactions followed by the antibody ($1\mu\text{g}/1 \times 10^6$ cells) for 1 h at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG2b ($1\mu\text{g}/1 \times 10^6$ cells) used under the same conditions. Acquisition of >10,000 events was performed.



1. Exosomes extracted from Hela cells
2. Exosomes extracted from Hela cells
3. Exosomes extracted from urine