




# PPA1 Monoclonal Antibody

|                            |  |
|----------------------------|--|
| <b>Product Code</b>        | CSB-MA614884A0m  |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| <b>Uniprot No.</b>         | Q15181   |
| <b>Immunogen</b>           | Recombinant Human Inorganic pyrophosphatase protein (1-289AA)  |
| <b>Raised In</b>           | Mouse  |
| <b>Species Reactivity</b>  | Human  |
| <b>Tested Applications</b> | ELISA, WB, IF, FC; Recommended dilution: WB: 1:1000-1:5000, IF: 1:50-1:200, FC: 1:50-1:200   |
| <b>Relevance</b>           | cytoplasm, cytosol, extracellular exosome, inorganic diphosphatase activity, diphosphate metabolic process, phosphate-containing compound metabolic process, tRNA aminoacylation for protein translation |
| <b>Form</b>                | Liquid   |
| <b>Conjugate</b>           | Non-conjugated   |
| <b>Storage Buffer</b>      | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, PH 7.4   |
| <b>Purification Method</b> | >95%, Protein G purified   |
| <b>Isotype</b>             | IgG1   |
| <b>Clonality</b>           | Monoclonal   |
| <b>Product Type</b>        | Monoclonal Antibody  |
| <b>Immunogen Species</b>   | Homo sapiens (Human)   |
| <b>Research Area</b>       | Cancer; Metabolism; Signal transduction  |
| <b>Gene Names</b>          | PPA1   |
| <b>Clone No.</b>           | 1H4E1  |

## Image



### Western Blot

Positive WB detected in: PPA1 antibody at 1:1000

Lane 1: HeLa whole cell lysate

Lane 2: HepG2 whole cell lysate

Lane 3: 293T whole cell lysate

Lane 4: MCF-7 whole cell lysate

Lane 5: U87 whole cell lysate


Secondary

Goat polyclonal to Mouse IgG at 1/20000 dilution


Predicted band size: 33KDa

Observed band size: 33 KDa


Exposure time: 5min



Immunofluorescence staining of Hela cells with CSB-MA614884A0m at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).



Overlay Peak curve showing HepG2 cells stained with CSB-MA614884A0m (red line) at 1:100. The cells were incubated in 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1µg/1\*10<sup>6</sup>cells) for 1h at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG1 (1µg/1\*10<sup>6</sup>cells) used under the same conditions. Acquisition of >10,000 events was performed.



Overlay Peak curve showing 293T cells stained with CSB-MA614884A0m (red line) at 1:100. The cells were incubated in 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1µg/1\*10<sup>6</sup>cells) for 1h at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG1 (1µg/1\*10<sup>6</sup>cells) used under the same conditions. Acquisition of >10,000 events was performed.