





# CCR8 Recombinant Monoclonal Antibody

<b>Product Code</b>	CSB-RA004847MA4HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	P51685
<b>Immunogen</b>	Recombinant Human CCR8 protein
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	FC
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	Affinity-chromatography
<b>Isotype</b>	hIgG1?lambda 2
<b>Clonality</b>	Monoclonal
<b>Product Type</b>	Recombinant Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Immunology
<b>Target Names</b>	CCR8
<b>Clone No.</b>	10A9

## Image



Untransfected HEK293T cells surface (green line) and transfected Human CCR8 HEK293T stable cells surface (red line) were stained with anti-CCR8 recombinant antibody ( $2\mu\text{g}/1 \times 10^6$  cells), washed and then followed by FITC-conjugated anti-Human IgG Fc antibody and analyzed with flow cytometry.



Untransfected CHO-K1 cells surface (green line) and transfected Human CCR8 CHO-K1 stable cells surface (red line) were stained with anti-CCR8 recombinant antibody ( $2\mu\text{g}/1 \times 10^6$  cells), washed and then followed by FITC-conjugated anti-Human IgG Fc antibody and analyzed with flow cytometry.

## Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.