





CDH17 Recombinant Monoclonal Antibody

Product Code	CSB-RA613267MA1HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q12864
Immunogen	Recombinant Human CDH17 protein
Species Reactivity	Human
Tested Applications	ELISA, FC; Recommended dilution: FC:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	Affinity-chromatography
Isotype	hIgG1
Clonality	Monoclonal
Product Type	Recombinant Antibody
Immunogen Species	Homo sapiens (Human)
Gene Names	CDH17
Clone No.	10B3

Image



The Binding Activity of Human CDH17 with Anti-CDH17 Recombinant Antibody
Activity: Measured by its binding ability in a functional ELISA. Immobilized Human CDH17 (CSB-MP613267HU) at 2 µg/mL can bind Anti-CDH17 recombinant antibody, the EC₅₀ is 3.095-4.451 ng/mL.



Untransfected HEK293T cells surface (green line) and transfected Human CDH17 HEK293T stable cells surface (red line) were stained with anti-CDH17 antibody (2µg/1*10⁶cells), washed and then followed by FITC-conjugated anti-Human IgG Fc antibody and analyzed with flow cytometry.



Description

The gene is isolated from the B cells produced by immunizing the animal with the recombinant human CDH17 protein and then cloned into a plasmid vector. The recombinant vector is introduced into host cell lines. The transfected cells are cultured to allow antibody expression. The CDH17 recombinant monoclonal antibody is purified from the cell lysate through affinity chromatography. It can be reacted with human CDH17 protein in the ELISA and FC applications.

CDH17 is primarily expressed in embryonic and adult intestinal epithelial cells, as well as in some pancreatic duct epithelial cells but is almost absent in liver cells, esophageal epithelial cells, and gastric mucosa in healthy individuals. Acting as a functional Ca^{2+} -dependent cell adhesion molecule, CDH17 exerts its adhesive function by directly interacting with the cellular cytoskeleton. Structural abnormalities and functional disruptions of CDH17 can lead to decreased cell-cell adhesion among tumor cells, resulting in loose tissue structure that is prone to detachment or deformation, thereby promoting tumor invasion and metastasis. Research has confirmed elevated expression levels of CDH17 in various human malignancies. Consequently, CDH17 can serve as a new objective indicator reflecting solid cancer's biological behavior and could potentially become a novel target for solid cancer treatment. Currently, clinical progress in CDH17-targeted drugs is limited, with most drugs in the preclinical stages. Therefore, the preparation of CDH17 protein could facilitate the development and research of CDH17-targeted clinical drugs.