





# INMT Recombinant Monoclonal Antibody

|                            |                                                                                                                               |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------|
| <b>Product Code</b>        | CSB-RA583078A0HU                                                                                                              |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.                                                                 |
| <b>Uniprot No.</b>         | O95050                                                                                                                        |
| <b>Immunogen</b>           | A synthesized peptide derived from Human INMT                                                                                 |
| <b>Species Reactivity</b>  | Human                                                                                                                         |
| <b>Tested Applications</b> | ELISA, IHC, FC; Recommended dilution: IHC:1:50-1:200, FC:1:50-1:200                                                           |
| <b>Form</b>                | Liquid                                                                                                                        |
| <b>Conjugate</b>           | Non-conjugated                                                                                                                |
| <b>Storage Buffer</b>      | Rabbit IgG in 10mM phosphate buffered saline , pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol. |
| <b>Purification Method</b> | Affinity-chromatography                                                                                                       |
| <b>Isotype</b>             | Rabbit IgG                                                                                                                    |
| <b>Clonality</b>           | Monoclonal                                                                                                                    |
| <b>Product Type</b>        | Recombinant Antibody                                                                                                          |
| <b>Immunogen Species</b>   | Homo sapiens (Human)                                                                                                          |
| <b>Research Area</b>       | Metabolism;Signal transduction                                                                                                |
| <b>Target Names</b>        | INMT                                                                                                                          |
| <b>Clone No.</b>           | 31G7                                                                                                                          |

## Image



IHC image of CSB-RA583078A0HU diluted at 1:50 and staining in paraffin-embedded human lung tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.24% DAB.



Overlay Peak curve showing SH-SY5Y cells stained with CSB-RA583078A0HU (red line) at 1:50. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1µg/1\*10<sup>6</sup>cells) for 45min at 4?. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4?. Control antibody (green line) was rabbit IgG (1µg/1\*10<sup>6</sup>cells) used under the same conditions. Acquisition of >10,000 events was performed.

**Usage**

For Research Use Only. Not for use in diagnostic or therapeutic procedures.