





# CD274 Recombinant Monoclonal Antibody

<b>Product Code</b>	CSB-RA878942MA1HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q9NZQ7
<b>Immunogen</b>	Recombinant Human CD274 protein
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, IHC, FC; Recommended dilution: IHC:1:20-1:200, FC:1:20-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	Affinity-chromatography
<b>Isotype</b>	Mouse IgG
<b>Clonality</b>	Monoclonal
<b>Product Type</b>	Recombinant Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Cancer;Immunology
<b>Gene Names</b>	CD274
<b>Clone No.</b>	24D6

## Image



IHC image of CSB-RA878942MA1HU diluted at 1:100 and staining in paraffin-embedded human lung cancer performed on a Leica Bond<sup>TM</sup> system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-Mouse IgG labeled by HRP and visualized using 0.05% DAB.



Overlay Peak curve showing Hela cells surface stained with CSB-RA878942MA1HU (red line) at 1:100. Then 10% normal goat serum was incubated to block non-specific protein-protein interactions followed by the antibody ( $1\mu\text{g}/1 \times 10^6$  cells) for 45 min at  $4^\circ\text{C}$ . The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/200 dilution for 35 min at  $4^\circ\text{C}$ . Isotype control antibody (green line) was mouse IgG1 ( $1\mu\text{g}/1 \times 10^6$  cells) used under the same conditions. Acquisition of  $>10,000$  events was performed.