





# BTLA Recombinant Monoclonal Antibody

<b>Product Code</b>	CSB-RA773799MA1HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q7Z6A9
<b>Immunogen</b>	Recombinant Human BTLA protein
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, FC; Recommended dilution: FC:1:50-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	Affinity-chromatography
<b>Isotype</b>	Human IgG4
<b>Clonality</b>	Monoclonal
<b>Product Type</b>	Recombinant Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Immunology;Stem cells
<b>Gene Names</b>	BTLA
<b>Clone No.</b>	10A10

## Image



The Binding Activity of Human BTLA with Anti-BTLA Recombinant Antibody  
Activity: Measured by its binding ability in a functional ELISA. Immobilized Human BTLA(CSB-MP773799HU1)at 2 µg/mL can bind Anti-BTLA recombinant antibody , the EC<sub>50</sub> is 11.56-13.00 ng/mL.



Untransfected Jurkat cells surface (green line) and transfected Human BTLA Jurkat stable cells surface (red line) were stained with anti-BTLA antibody (2µg/1\*10<sup>6</sup>cells), washed and then followed by APC-conjugated anti-Human IgG Fc antibody and analyzed with flow cytometry.

