





# MUC13 Recombinant Monoclonal Antibody

<b>Product Code</b>	CSB-RA887973MA1HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Immunogen</b>	Recombinant Human MUC13 protein
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA; Recommended dilution: ELISA:1:5000-1:50000
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	Affinity-chromatography
<b>Isotype</b>	hIgG1
<b>Clonality</b>	Monoclonal
<b>Product Type</b>	Recombinant Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Purity</b>	Greater than 95% as determined by SEC-HPLC.
<b>Research Area</b>	Cancer;Signal transduction
<b>Target Names</b>	MUC13
<b>Clone No.</b>	6H10


## Image




The Binding Activity of Human MUC13 with Anti-MUC13 recombinant antibody  
Activity: Measured by its binding ability in a functional ELISA. Immobilized Human MUC13 (CSB-MP887973HU) at 2 µg/mL can bind Anti-MUC13 recombinant antibody. The EC<sub>50</sub> is 3.806-5.269 ng/mL.



The Binding Activity of Human MUC13 with Anti-MUC13 recombinant antibody  
Activity: Measured by its binding ability in a functional ELISA. Immobilized Human MUC13 (CSB-YP887973HU) at 2 µg/mL can bind Anti-MUC13 recombinant antibody. The EC<sub>50</sub> is 2.592-2.923 ng/mL.



The purity of MUC13 was greater than 95% as determined by SEC-HPLC



The purity of MUC13 was greater than 95% as determined by SEC-HPLC

## Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.