





MMAE&MMAF Recombinant Monoclonal Antibody

Product Code	CSB-RA90389011m
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Immunogen	MMAE-BSA
Species Reactivity	Chemical
Tested Applications	ELISA; Recommended dilution: ELISA:1:5000-1:50000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	Affinity-chromatography
Isotype	IgG2b
Clonality	Monoclonal
Product Type	Recombinant Antibody
Purity	Greater than 95% as determined by SEC-HPLC.
Research Area	Others
Target Names	MMAE; MMAF
Clone No.	25G12B5


Image



The Binding Activity of Disitamab vedotin (Synonyms: RC48) with Anti-MMAE&MMAF antibody
Activity: Measured by its binding ability in a functional ELISA. Immobilized Disitamab vedotin (Synonyms: RC48) at 2 µg/mL can bind Anti-MMAE&MMAF antibody, the EC₅₀ is 3.523 to 5.356 ng/mL.




The Binding Activity of Depatuxizumab mafodotin (Synonyms: ABT-414) with Anti-MMAE&MMAF antibody
Activity: Measured by its binding ability in a functional ELISA. Immobilized Depatuxizumab mafodotin (Synonyms: ABT-414) at 2 µg/mL can bind Anti-MMAE&MMAF antibody, the EC₅₀ is 3.022 to 4.198 ng/mL.



The Binding Activity of Disitamab vedotin (Synonyms: RC48) and Depatuxizumab mafodotin (Synonyms: ABT-414) with Anti-MMAE&MMAF antibody

Activity: Measured by its binding ability in a functional ELISA. Immobilized Disitamab vedotin (Synonyms: RC48) and Depatuxizumab mafodotin (Synonyms: ABT-414) at 2 μg/mL can bind Anti-MMAE&MMAF antibody, the EC₅₀ is 3.523 to 5.356 ng/mL and 3.022 to 4.198 ng/mL, respectively.



The purity of MMAE&MMAF was greater than 95% as determined by SEC-HPLC

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.