





# CD79B Recombinant Monoclonal Antibody

<b>Product Code</b>	CSB-RA004958MA2HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	P40259
<b>Immunogen</b>	Recombinant Human CD79B protein
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, FC; Recommended dilution: ELISA:1:5000-1:50000, FC:1:50-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	Affinity-chromatography
<b>Isotype</b>	hIgG1
<b>Clonality</b>	Monoclonal
<b>Product Type</b>	Recombinant Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Immunology
<b>Target Names</b>	CD79B
<b>Clone No.</b>	8E9


## Image




The Binding Activity of Human CD79B with Anti-CD79B recombinant antibody  
Activity: Measured by its binding ability in a functional ELISA. Immobilized Human CD79A(CSB-MP004958HU1) at 2 µg/mL can bind Anti-CD79B recombinant antibody. The EC<sub>50</sub> is 3.214-3.642 ng/mL.




The Binding Activity of Human CD79A&CD79B with Anti-CD79B recombinant antibody  
Activity: Measured by its binding ability in a functional ELISA. Immobilized Human CD79A&CD79B(CSB-MP5861HU) at 2 µg/mL can bind Anti-CD79B recombinant antibody. The EC<sub>50</sub> is 14.85-17.47 ng/mL.



The Binding Activity of Human CD79B with Anti-CD79B recombinant antibody  
 Activity: Measured by its binding ability in a functional ELISA. Immobilized Human CD79B(CSB-MP004958HU1p7) at 2 µg/mL can bind Anti-CD79B recombinant antibody. The EC<sub>50</sub> is 31.97-35.69 ng/mL.



Untransfected HEK-293T cells (green line) and transfected Human CD79B HEK-293T stable cells (red line) were stained with anti-CD79B antibody(CSB-RA004958MA2HU) (2µg/1\*10<sup>6</sup>cells), washed and then followed by APC-conjugated anti-Human IgG Fc antibody and analyzed with flow cytometry.



Untransfected HEK-293T cells (green line) and transfected Human CD79A&CD79B HEK-293T stable cells (red line) were stained with anti-CD79B antibody(CSB-RA004958MA2HU) (2µg/1\*10<sup>6</sup>cells), washed and then followed by APC-conjugated anti-Human IgG Fc antibody and analyzed with flow cytometry.

**Usage**

For Research Use Only. Not for use in diagnostic or therapeutic procedures.