





MICA/B Recombinant Monoclonal Antibody

Product Code	CSB-RA56059MA1HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	J9TPG6
Immunogen	Recombinant Human MICA/B protein
Species Reactivity	Human, Macaca fascicularis
Tested Applications	ELISA; Recommended dilution: ELISA:1:5000-1:50000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	Affinity-chromatography
Isotype	hIgG1
Clonality	Monoclonal
Product Type	Recombinant Antibody
Immunogen Species	Homo sapiens (Human)
Research Area	Others
Target Names	MICA/B
Clone No.	5B8


Image




The Binding Activity of Human MICB with Anti-MICA/B Recombinant Antibody
Activity: Measured by its binding ability in a functional ELISA. Immobilized Human MICB (CSB-MP6145HU) at 2 µg/mL can bind Anti-MICA/B recombinant antibody. The EC₅₀ is 1.087-1.369 ng/mL.




The Binding Activity of Human MICA with Anti-MICA/B recombinant antibody
Activity: Measured by its binding ability in a functional ELISA. Immobilized Human MICA (CSB-MP5606HU) at 2 µg/mL can bind Anti-MICA/B recombinant antibody. The EC₅₀ is 0.9163-1.108 ng/mL.




The Binding Activity of Macaca fascicularis MICB with Anti-MICA/B recombinant antibody Activity: Measured by its binding ability in a functional ELISA. Immobilized Macaca fascicularis MICB(CSB-MP6584MOV(A4) at 2 μ g/mL can bind Anti-MICA/B recombinant antibody. The EC_{50} is 1.578-1.831 ng/mL.



The Binding Activity of Human MICB with Anti-MICA/B recombinant antibody Activity: Measured by its binding ability in a functional ELISA. Immobilized Human MICB(CSB-MP5608HU) at 2 μ g/mL can bind Anti-MICA/B recombinant antibody. The EC_{50} is 0.9180-1.062 ng/mL.



The Binding Activity of Human MICB with Anti-MICA/B recombinant antibody Activity: Measured by its binding ability in a functional ELISA. Immobilized Human MICB(CSB-MP5607HU1) at 2 μ g/mL can bind Anti-MICA/B recombinant antibody. The EC_{50} is 26.77-32.93 ng/mL.



The Binding Activity of Human MICA with Anti-MICA/B recombinant antibody Activity: Measured by its binding ability in a functional ELISA. Immobilized Human MICA(CSB-MP5605HU) at 2 μ g/mL can bind Anti-MICA/B recombinant antibody. The EC_{50} is 0.6924-0.9195 ng/mL.

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.