





# CCL17 Recombinant Monoclonal Antibody

<b>Product Code</b>	CSB-RA856406MA1HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q92583
<b>Immunogen</b>	Recombinant Human CCL17 protein
<b>Species Reactivity</b>	Human, Mouse, Macaca fascicularis
<b>Tested Applications</b>	ELISA, FC; Recommended dilution: ELISA:1:5000-1:50000, FC:1:50-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	Affinity-chromatography
<b>Isotype</b>	hIgG2
<b>Clonality</b>	Monoclonal
<b>Product Type</b>	Recombinant Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Immunology
<b>Target Names</b>	CCL17
<b>Clone No.</b>	7D3


## Image




The Binding Activity of Human CCL17 with Anti-CCL17 recombinant antibody  
Activity: Measured by its binding ability in a functional ELISA. Immobilized Anti-CCL17 recombinant antibody at 2 µg/mL can bind Human CCL17 protein (CSB-MP856406HU) . The EC<sub>50</sub> is 2.403-2.741 ng/mL.



The Binding Activity of Macaca mulatta CCL17 with Anti-CCL17 recombinant antibody  
Activity: Measured by its binding ability in a functional ELISA. Immobilized Anti-CCL17 recombinant antibody at 2 µg/mL can bind Mouse Ccl17 protein (CSB-MP6512MO). The EC<sub>50</sub> is 0.9792-1.486 ng/mL.



The Binding Activity of Macaca mulatta CCL17 with Anti-CCL17 recombinant antibody Activity: Measured by its binding ability in a functional ELISA. Immobilized Anti-CCL17 recombinant antibody at 2 µg/mL can bind Macaca mulatta CCL17 protein (CSB-MP811562MOW). The EC<sub>50</sub> is 1.953-2.469 ng/mL.



Overlay Peak curve showing Raji cells stained with CSB-RA856406MA1HU (red line) at 1:100. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100 for 10min. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1\*10<sup>6</sup>cells) for 45min at 4?. The secondary antibody used was Fluorescein (FITC) AffiniPure Goat Anti-Human IgG, Fcγ fragment specific at 1:200 dilution for 35 min at 4?. Control antibody (green line) was human IgG1 (1ug/1\*10<sup>6</sup>cells) used under the same conditions. Acquisition of >10,000 events was performed.

**Usage**

For Research Use Only. Not for use in diagnostic or therapeutic procedures.