




EEF1A1 Recombinant Monoclonal Antibody

Product Code	CSB-RA019747A0HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P68104
Immunogen	A synthesized peptide derived from human EEF1A1
Species Reactivity	Human, Mouse, Rat
Tested Applications	ELISA, WB, IF, FC; Recommended dilution: WB:1:500-1:2000, IF:1:50-1:200, FC:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Rabbit IgG in 10mM phosphate buffered saline , pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Purification Method	Affinity-chromatography
Isotype	Rabbit IgG
Clonality	Monoclonal
Product Type	Recombinant Antibody
Immunogen Species	Homo sapiens (Human)
Target Names	EEF1A1
Clone No.	9C12

Image



Western Blot

Positive WB detected in: HepG2 whole cell lysate(30µg),U-937 whole cell lysate(30µg), HL-60 whole cell lysate(30µg), HeLa whole cell lysate(30µg), PC-3 whole cell lysate(30µg),THP-1 whole cell lysate(30µg),NIH/3T3 whole cell lysate(30µg),HEK293 whole cell lysate(30µg), Rat brain tissue lysate(30µg)

All lanes: eEF1A1 antibody at 1:1000


Secondary

Goat polyclonal to rabbit IgG at 1/40000 dilution


Predicted band size: 50 kDa

Observed band size: 55 kDa

Exposure time?20s



Immunofluorescence staining of HepG2 cell with CSB-RA019747A0HU at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Overlay Peak curve showing jurkat cells stained with CSB-RA019747A0HU (red line) at 1:100. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100 for 10min. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 45min at 4?. The secondary antibody used was FITC-conjugated goat anti-rabbit IgG (H+L) at 1/200 dilution for 35min at 4?. Control antibody (green line) was Rabbit IgG (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.