




IDH1 Recombinant Monoclonal Antibody

Product Code	CSB-RA916901A0HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	O75874
Immunogen	A synthesized peptide derived from Human IDH1 protein
Species Reactivity	Human, Mouse
Tested Applications	ELISA, WB, FC; Recommended dilution: WB:1:500-1:5000, FC:1:20-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Rabbit IgG in 10mM phosphate buffered saline , pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Purification Method	Affinity-chromatography
Isotype	Rabbit IgG
Clonality	Monoclonal
Product Type	Recombinant Antibody
Immunogen Species	Homo sapiens (Human)
Target Names	IDH1
Clone No.	4H3

Image



Western Blot

Positive WB detected in: HepG2 whole cell lysate(30μg), COLO205 whole cell lysate(30μg), HEK293 whole cell lysate(30μg), HeLa whole cell lysate(30μg), NIH/3T3 whole cell lysate(30μg), A431 whole cell lysate(30μg), Mouse Brain tissue whole cell lysate(30μg)

All lanes: IDH1 antibody at 1:1000


Secondary

Goat polyclonal to rabbit IgG at 1/40000 dilution

Predicted band size: 47 kDa

Observed band size: 45 kDa

Exposure time?30S



Overlay Peak curve showing HepG2 cells stained with CSB-RA916901A0HU (red line) at 1:100. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100 for 10min. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 45min at 4?. The secondary antibody used was FITC-conjugated goat anti-rabbit IgG (H+L) at 1/200 dilution for 35min at 4?. Control antibody (green line) was Rabbit IgG (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.