



Recombinant Rat Dual specificity mitogen-activated protein kinase kinase 1 (Map2k1)

Product Code	CSB-EP013409RA-B
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	Q01986
Product Type	Recombinant Protein
Immunogen Species	Rattus norvegicus (Rat)
Purity	>85% (SDS-PAGE)
Sequence	PKKKPTPIQ LNPAPDGS AV NGTSSAETNL EALQKKLEEL ELDEQQRKRL EAFLTQKQKV GELKDDDFEK ISELGAGNGG VVFKVSHKPS GLVMARKLIH LEIKPAIRNQ IIRELQVLHE CNSPIYVGFY GAFYSDGEIS ICMEHMDGGS LDQVLKAGR IPEQILGKVS IAVIKGLTYL REKHKIMHRD VKPSNILVNS RGEIKLDFG VSGQLIDSMA NSFVGT RSYM SPERLQGTHY SVQSDIWSMG LSLVEMAVGR YPIPPDAKE LELLFGCQVE GDA AETPPRP RTPGRPLSSY GMDSRPPMAI FELLDYIVNE PPPKLPSGVF SLEFQDFV NK CLIKNPAERA DLKQLMVHAF IKRSDAEEVD FAGWLCSTIG LNQPSTPTHA ASI
Source	E.coli
Target Names	Map2k1
Protein Names	Recommended name: Dual specificity mitogen-activated protein kinase kinase 1 Short name= MAP kinase kinase 1 Short name= MAPKK 1 EC= 2.7.12.2 Alternative name(s): ERK activator kinase 1 MAPK/ERK kinase 1 Short name=
Expression Region	2-393
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	This protein is a member of the dual specificity protein kinase family, which acts as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein kinase lies upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon wide variety of extra- and intracellular signals. As an essential component of MAP kinase signal transduction pathway, this kinase is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.



Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.