



Recombinant Rat Mitogen-activated protein kinase 8 (Mapk8)

Product Code	CSB-YP013466RA
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P49185
Product Type	Recombinant Protein
Immunogen Species	Rattus norvegicus (Rat)
Purity	>85% (SDS-PAGE)
Sequence	MSRSKRDNF YSVEIADSTF TVLKRYQNLK PIGSGAQQIV CAAYDAILER NVAIKKLSRP FQNQTHAKRA YRELVLMKCV NHKNIIGLLN VFTPQKSLEE FQDVYIVMEL MDANLCQVIQ MELDHERMSY LLYQMLCGIK HLHSAGIIHR DLKPSNIVVK SDCTLKILDF GLARTAGTSF MTPYVVTRY YRAPEVILGM GYKENVDLWS VGCIMGEMVC LKILFPGRDY IDQWNKVIEQ LGTPCPEFMK KLQPTVRTYV ENRPKYAGYS FEKLFPDVLV PADSEHNKLN ASQARDLLSK MLVIDASKRI SVDEALQHPY INVWYDPSEA EAPPPKIPDK QLDEREHTIE EWKELIYKEV MDLEERTKNG VIRGQPSPLG AAVINGSQHP VSSPSVNDMS SMSTDPTLAS D
Source	Yeast
Target Names	Mapk8
Protein Names	Recommended name: Mitogen-activated protein kinase 8 Short name= MAP kinase 8 Short name= MAPK 8 EC= 2.7.11.24 Alternative name(s): SAPK gamma Stress-activated protein kinase JNK1 c-Jun N-terminal kinase 1 p54 gamm
Expression Region	1-411
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full length protein
Target Details	This protein is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various cell stimuli, and targets specific transcription factors, and thus mediates immediate-early gene expression in response to cell stimuli. The activation of this kinase by tumor-necrosis factor alpha (TNF-alpha) is found to be required for TNF-alpha induced apoptosis. This kinase is also involved in UV radiation induced apoptosis, which is thought to be related to cytochrom c-mediated cell death pathway. Studies of the mouse counterpart of this gene suggested that this kinase play a key role in T cell proliferation, apoptosis and differentiation. Four alternatively spliced



transcript variants encoding distinct isoforms have been reported.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.