



Recombinant Rat Mitogen-activated protein kinase 9 (Mapk9)

Product Code	CSB-EP013471RA
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P49186
Product Type	Recombinant Protein
Immunogen Species	Rattus norvegicus (Rat)
Purity	>85% (SDS-PAGE)
Sequence	MSDSKSDGQF YSVQVADSTF TVLKRYQQLK PIGSGAQQGIV CAAFDTV LGI NVAVK KLSRP FQNQTHAKRA YRELVLLKCV NHKNIISLLN VFTPQK TLEE FQDVYLV MEL MDANLCQVIH MELDHERMSY LLYQMLCGIK HLHSAGIIHR DLKPSNIVVK SDCTLKILDF GLARTACTNF MTPYVVTRY YRAPEVILGM GYKENVDIWS VGCIMGELVK GCVIFQGDH IDQWNKVIEQ LGTPSAEFMK KLQPTVRNYV ENRPKYPGIK FEELFPDWIF PSESERDKIK TSQARDLLSK MLVIDPDKRI SVDEALRHPY ITVWYDPAEA EAPPPQIYDA QLEEREHAIE EWKELIYKEV MDWEERSKNG VKDQPSDAAV SSKATPSQSS SINDISSMST EHTLASDTDS SLDASTGPLE GCR
Source	E.coli
Target Names	Mapk9
Protein Names	Recommended name: Mitogen-activated protein kinase 9 Short name= MAP kinase 9 Short name= MAPK 9 EC= 2.7.11.24 Alternative name(s): SAPK-alpha Stress-activated protein kinase JNK2 c-Jun N-terminal kinase 2 p54-alpha
Expression Region	1-423
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full length protein
Target Details	This protein is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase targets specific transcription factors, and thus mediates immediate-early gene expression in response to various cell stimuli. It is most closely related to MAPK8, both of which are involved in UV radiation induced apoptosis, thought to be related to the cytochrome c-mediated cell death pathway. This gene and MAPK8 are also known as c-Jun N-terminal kinases. This kinase blocks the ubiquitination of tumor suppressor p53, and thus it increases the stability of p53 in nonstressed cells. Studies of this gene's mouse counterpart suggest a key role in T-cell differentiation. Several alternatively spliced transcript variants encoding distinct isoforms have been



reported.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.