



Recombinant Mouse Mitogen-activated protein kinase 14 (Mapk14)

Product Code	CSB-YP013453MO
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P47811
Product Type	Recombinant Protein
Immunogen Species	Mus musculus (Mouse)
Purity	>85% (SDS-PAGE)
Sequence	SQERPTFYR QELNKTIWEV PERYQNLSPV GSGAYGSVCA AFDTKTGHRV AVKKLSRPFQ SIIHAKRTYR ELRLLKHKH ENVIGLLDVF TPARSLEEFN DVYLVTHLMG ADLNNIVKCQ KLTDDHVQFL IYQILRGLKY IHSADIIHRD LKPSNLAVNE DCELKILDFG LARHTDDEMT GYVATRWYRA PEIMLNWMHY NQTVDIWSVG CIMAELLTGR TLPFGTDHID QLKLILRLVG TPGAELLKKI SSESARNYIQ SLAQMPKMNF ANVFIGANPL AVDLLEKMLV LDSDKRITAA QALAHAYFAQ YHDPDDEPVA DPYDQSFESR DLLIDEWKSL TYDEVISFVP PPLDQEEMES
Source	Yeast
Target Names	Mapk14
Protein Names	Recommended name: Mitogen-activated protein kinase 14 Short name= MAP kinase 14 Short name= MAPK 14 EC= 2.7.11.24 Alternative name(s): CRK1 Mitogen-activated protein kinase p38 alpha Short name= MAP kinase p38 alpha
Expression Region	2-360
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	This protein is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the



contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.