

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

CYTH3 Antibody

CSB-PA006469GA01HU Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Lippon receipt store at -20°C or -80°C. Avoid repeated freeze
O43739
Human CYTH3
Rabbit
Human,Mouse,Rat
ELISA,WB
PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.
Antigen Affinity purified
IgG
cytohesin 3;CYTH3;ARNO3;GRP1;PSCD3 ;
Purified Rabbit Anti human PolyClonal Antibody
Homo sapiens (Human)
CYTH3
This gene encodes a member of the PSCD (pleckstrin homology, Sec7 and coiled-coil domains) family. PSCD family members have identical structural organization that consists of an N-terminal coiled-coil motif, a central Sec7 domain, and a C-terminal pleckstrin homology (PH) domain. The coiled-coil motif is involved in homodimerization, the Sec7 domain contains guanine-nucleotide exchange protein (GEP) activity, and the PH domain interacts with phospholipids and is responsible for association of PSCDs with membranes. Members of this family appear to mediate the regulation of protein sorting and membrane trafficking. This encoded protein is involved in the control of Golgi structure and function, and it may have a physiological role in regulating ADP-ribosylation factor protein 6 (ARF) functions, in addition to acting on ARF1.

1