

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

AGO2 Antibody

Product CodeCSB-PA007520GA01HUStorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.Q9UKV8ImmunogenHuman EIF2C2Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoi cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaseukaryotic translation initiation factor 2C, 2;EIF2C2;AGO2;MGC3Product TypeHomo sapiens (Human)	
Uniprot No.Q9UKV8ImmunogenHuman EIF2C2Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoi cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaseukaryotic translation initiation factor 2C, 2;EIF2C2;AGO2;MGC3Product TypePurified Rabbit Anti human PolyClonal Antibody	
ImmunogenHuman EIF2C2Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoi cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaseukaryotic translation initiation factor 2C, 2;EIF2C2;AGO2;MGC3Product TypePurified Rabbit Anti human PolyClonal Antibody	
Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoi cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaseukaryotic translation initiation factor 2C, 2;EIF2C2;AGO2;MGC3Product TypePurified Rabbit Anti human PolyClonal Antibody	
Species ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoi cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaseukaryotic translation initiation factor 2C, 2;EIF2C2;AGO2;MGC3Product TypePurified Rabbit Anti human PolyClonal Antibody	
Tested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoi cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaseukaryotic translation initiation factor 2C, 2;EIF2C2;AGO2;MGC3Product TypePurified Rabbit Anti human PolyClonal Antibody	
Storage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoi cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaseukaryotic translation initiation factor 2C, 2;EIF2C2;AGO2;MGC3Product TypePurified Rabbit Anti human PolyClonal Antibody	
cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaseukaryotic translation initiation factor 2C, 2;EIF2C2;AGO2;MGC3Product TypePurified Rabbit Anti human PolyClonal Antibody	
IsotypeIgGAliaseukaryotic translation initiation factor 2C, 2;EIF2C2;AGO2;MGC3Product TypePurified Rabbit Anti human PolyClonal Antibody	d freeze / thaw
Aliaseukaryotic translation initiation factor 2C, 2;EIF2C2;AGO2;MGC3Product TypePurified Rabbit Anti human PolyClonal Antibody	
Product Type Purified Rabbit Anti human PolyClonal Antibody	
	183;Q10 ;
Immunogen Species Homo sapiens (Human)	
Target Names AGO2	
Target DetailsThis gene encodes a member of the Argonaute family of proteins role in RNA interference. The encoded protein is highly basic, and PAZ domain and a PIWI domain. It may interact with dicer1 and p short-interfering-RNA-mediated gene silencing. Multiple transcrip encoding different isoforms have been found for this gene.	d contains a play a role in

1