



# FXVD1 Antibody

<b>Product Code</b>	CSB-PA009089GA01HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	O00168
<b>Immunogen</b>	Human FXVD1
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human,Mouse,Rat
<b>Tested Applications</b>	ELISA,WB
<b>Storage Buffer</b>	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3. -20°C, Avoid freeze / thaw cycles.
<b>Purification Method</b>	Antigen Affinity purified
<b>Isotype</b>	IgG
<b>Alias</b>	FXVD domain containing ion transport regulator 1;FXVD1;MGC44983;PLM ;Phospholemman
<b>Product Type</b>	Purified Rabbit Anti human PolyClonal Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Target Names</b>	FXVD1
<b>Target Details</b>	<p>This gene encodes a member of a family of small membrane proteins that share a 35-amino acid signature sequence domain, beginning with the sequence PFXVD and containing 7 invariant and 6 highly conserved amino acids. The approved human gene nomenclature for the family is FXVD-domain containing ion transport regulator. Mouse FXVD5 has been termed RIC (Related to Ion Channel). FXVD2, also known as the gamma subunit of the Na,K-ATPase, regulates the properties of that enzyme. FXVD1 (phospholemman), FXVD2 (gamma), FXVD3 (MAT-8), FXVD4 (CHIF), and FXVD5 (RIC) have been shown to induce channel activity in experimental expression systems. Transmembrane topology has been established for two family members (FXVD1 and FXVD2), with the N-terminus extracellular and the C-terminus on the cytoplasmic side of the membrane. This protein is a plasma membrane substrate for several kinases, including protein kinase A, protein kinase C, NIMA kinase, and myotonic dystrophy kinase. It is thought to form an ion channel or regulate ion channel activity. Transcript variants with different 5 UTR sequences have been described in the literature.</p>
<b>Usage</b>	For Research Use Only. Not for use in diagnostic or therapeutic procedures.