

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

LAT Antibody

cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslinker for activation of T cells;LAT;LAT1;pp36 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This		
Uniprot No.O43561ImmunogenHuman LATRaised InRabbitSpecies ReactivityHumanTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslinker for activation of T cells;LAT;LAT1;pp36 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Product Code	CSB-PA012767GA01HU
ImmunogenHuman LATRaised InRabbitSpecies ReactivityHumanTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslinker for activation of T cells;LAT;LAT1;pp36 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitRaised InRabbitSpecies ReactivityHumanTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslinker for activation of T cells;LAT;LAT1;pp36 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Uniprot No.	O43561
Species ReactivityHumanTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslinker for activation of T cells;LAT;LAT1;pp36 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Immunogen	Human LAT
Tested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslinker for activation of T cells;LAT;LAT1;pp36 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Raised In	Rabbit
Storage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslinker for activation of T cells;LAT;LAT1;pp36 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Species Reactivity	Human
cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslinker for activation of T cells;LAT;LAT1;pp36 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Tested Applications	ELISA,WB
IsotypeIgGAliaslinker for activation of T cells;LAT;LAT1;pp36 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Storage Buffer	PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.
Aliaslinker for activation of T cells;LAT;LAT1;pp36 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Purification Method	Antigen Affinity Purified
Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Isotype	lgG
Immunogen SpeciesHomo sapiens (Human)Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Alias	linker for activation of T cells;LAT;LAT1;pp36 ;
Target NamesLATTarget DetailsThis protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Product Type	Purified Rabbit Anti human PolyClonal Antibody
Target Details This protein is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Immunogen Species	Homo sapiens (Human)
activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative	Target Names	LAT
	Target Details	adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative

1