

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

## MAP2K6 Antibody

<ul> <li>CSB-PA013415GA01HU</li> <li>Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.</li> <li>P52564</li> <li>Human MAP2K6</li> <li>Rabbit</li> <li>Human,Mouse,Rat</li> <li>ELISA,WB,IHC</li> <li>PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.</li> <li>Antigen Affinity Purified</li> <li>IgG</li> <li>mitogen-activated protein kinase kinase</li> </ul>
<ul> <li>P52564</li> <li>Human MAP2K6</li> <li>Rabbit</li> <li>Human,Mouse,Rat</li> <li>ELISA,WB,IHC</li> <li>PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.</li> <li>Antigen Affinity Purified</li> <li>IgG</li> </ul>
<ul> <li>Human MAP2K6</li> <li>Rabbit</li> <li>Human,Mouse,Rat</li> <li>ELISA,WB,IHC</li> <li>PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.</li> <li>Antigen Affinity Purified</li> <li>IgG</li> </ul>
Rabbit         Human,Mouse,Rat         ELISA,WB,IHC         PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.         Antigen Affinity Purified         IgG
Human,Mouse,Rat         ELISA,WB,IHC         PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.         Antigen Affinity Purified         IgG
<ul> <li>ELISA,WB,IHC</li> <li>PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.</li> <li>Antigen Affinity Purified</li> <li>IgG</li> </ul>
<ul> <li>PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.</li> <li>Antigen Affinity Purified</li> <li>IgG</li> </ul>
cycles. Antigen Affinity Purified IgG
IgG
mitogen-activated protein kinase kinase
6;MAP2K6;MAPKK6;MEK6;MKK6;PRKMK6;SAPKK3 ;
Purified Rabbit Anti human PolyClonal Antibody
Homo sapiens (Human)
MAP2K6
This gene encodes a member of the dual specificity protein kinase family, which functions as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein phosphorylates and activates p38 MAP kinase in response to inflammatory cytokines or environmental stress. As an essential component of p38 MAP kinase mediated signal transduction pathway, this gene is involved in many cellular processes such as stress induced cell cycle arrest, transcription activation and apoptosis.

1