

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

MAPK11 Antibody

 CSB-PA013450GA01HU Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Q15759 Human MAPK11 Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.
Q15759 Human MAPK11 Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
Human MAPK11 Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
Antigen Affinity purified
IgG
mitogen-activated protein kinase 11;MAPK11;P38B;P38BETA2;PRKM11;SAPK2;SAPK2B;p38-2;p38Beta ;
Purified Rabbit Anti human PolyClonal Antibody
Homo sapiens (Human)
MAPK11
This protein is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation, and development. This kinase is most closely related to p38 MAP kinase, both of which can be activated by proinflammatory cytokines and environmental stress. This kinase is activated through its phosphorylation by MAP kinase kinases (MKKs), preferably by MKK6. Transcription factor ATF2/CREB2 has been shown to be a substrate of this kinase.

1