

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

MGAT2 Antibody

 CSB-PA013774GA01HU Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Q10469 Human MGAT2 Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity purified
Q10469 Human MGAT2 Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity purified
Human MGAT2 Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity purified
Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity purified
Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity purified
ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity purified
PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity purified
cycles. Antigen Affinity purified
IgG
mannosyl (alpha-1,6-)-glycoprotein beta-1,2-N- acetylglucosaminyltransferase;MGAT2;CDGS2;GLCNACTII;GNT-II;GNT2;
Purified Rabbit Anti human PolyClonal Antibody
Homo sapiens (Human)
MGAT2
The product of this gene is a Golgi enzyme catalyzing an essential step in the conversion of oligomannose to complex N-glycans. The enzyme has the typical glycosyltransferase domains: a short N-terminal cytoplasmic domain, a hydrophobic non-cleavable signal-anchor domain, and a C-terminal catalytic domain. Mutations in this gene may lead to carbohydrate-deficient glycoprotein syndrome, type II. The coding region of this gene is intronless. Transcript variants with a spliced 5 UTR may exist, but their biological validity has not been determined.

1