

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

NAGA Antibody

Product CodeCSB-PA015414GA01HUStorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.P17050ImmunogenHuman NAGARaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasN-acetylgalactosaminidase, alpha-;NAGA;D22S674;GALB ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesNAGATarget DetailsNAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminidase, store s		
Uniprot No.P17050ImmunogenHuman NAGARaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasN-acetylgalactosaminidase, alpha-;NAGA;D22S674;GALB ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesNAGANAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Product Code	CSB-PA015414GA01HU
ImmunogenHuman NAGARaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasN-acetylgalactosaminidase, alpha-;NAGA;D22S674;GALB ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesNAGATarget DetailsNAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasN-acetylgalactosaminidase, alpha-;NAGA;D22S674;GALB ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesNAGATarget DetailsNAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Uniprot No.	P17050
Species ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasN-acetylgalactosaminidase, alpha-;NAGA;D22S674;GALB ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesNAGATarget DetailsNAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Immunogen	Human NAGA
Tested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasN-acetylgalactosaminidase, alpha-;NAGA;D22S674;GALB ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesNAGATarget DetailsNAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Raised In	Rabbit
Storage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasN-acetylgalactosaminidase, alpha-;NAGA;D22S674;GALB ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesNAGATarget DetailsNAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Species Reactivity	Human,Mouse,Rat
thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasN-acetylgalactosaminidase, alpha-;NAGA;D22S674;GALB ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesNAGANAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Tested Applications	ELISA,WB,IHC
IsotypeIgGAliasN-acetylgalactosaminidase, alpha-;NAGA;D22S674;GALB ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesNAGATarget DetailsNAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Storage Buffer	
AliasN-acetylgalactosaminidase, alpha-;NAGA;D22S674;GALB ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesNAGATarget DetailsNAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Purification Method	Antigen Affinity purified
Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesNAGATarget DetailsNAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Isotype	IgG
Immunogen SpeciesHomo sapiens (Human)Target NamesNAGATarget DetailsNAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Alias	N-acetylgalactosaminidase, alpha-;NAGA;D22S674;GALB;
Target Names NAGA Target Details NAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Product Type	Purified Rabbit Anti human PolyClonal Antibody
Target Details NAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Immunogen Species	Homo sapiens (Human)
cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II	Target Names	NAGA
	Target Details	cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II

1