



# PDCD10 Antibody

|                            |  |
|----------------------------|--|
| <b>Product Code</b>        | CSB-PA017665GA01HU   |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| <b>Uniprot No.</b>         | Q9BUL8   |
| <b>Immunogen</b>           | Human PDCD10   |
| <b>Raised In</b>           | Rabbit   |
| <b>Species Reactivity</b>  | Human,Mouse,Rat  |
| <b>Tested Applications</b> | ELISA,WB,IHC   |
| <b>Storage Buffer</b>      | PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.3. -20°C, Avoid freeze / thaw cycles.   |
| <b>Purification Method</b> | Antigen Affinity Purified  |
| <b>Isotype</b>             | IgG  |
| <b>Alias</b>               | programmed cell death 10;PDCD10;CCM3;MGC1212;MGC24477;TFAR15 ;   |
| <b>Product Type</b>        | Purified Rabbit Anti human PolyClonal Antibody   |
| <b>Immunogen Species</b>   | Homo sapiens (Human)   |
| <b>Target Names</b>        | PDCD10   |
| <b>Target Details</b>      | <p>This gene encodes an evolutionarily conserved protein associated with cell apoptosis. The protein interacts with the serine/threonine protein kinase MST4 to modulate the extracellular signal-regulated kinase (ERK) pathway. It also interacts with and is phosphorylated by serine/threonine kinase 25, and is thought to function in a signaling pathway essential for vascular development. Mutations in this gene are one cause of cerebral cavernous malformations, which are vascular malformations that cause seizures and cerebral hemorrhages. Multiple alternatively spliced variants, encoding the same protein, have been identified.</p> |
| <b>Usage</b>               | For Research Use Only. Not for use in diagnostic or therapeutic procedures.  |