

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

## PIK3CA Antibody

Product CodeCSB-PA017998GA01HUStorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.P42336ImmunogenHuman PIK3CARaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasphosphoinositide-3-kinase, catalytic, alpha polypeptide;PIK3CA;MGC142161;MGC142163;PI3K;p110-alpha ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesPIK3CATarget NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has been inpulcated in cervical cancers.		
Uniprot No.P42336UmmunogenHuman PIK3CARaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasphosphoinositide-3-kinase, catalytic, alpha polypeptide;PIK3CA;MGC142161;MGC142163;PI3K;p110-alpha ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Product Code	CSB-PA017998GA01HU
ImmunogenHuman PIK3CARaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasphosphoinositide-3-kinase, catalytic, alpha polypeptide;PIK3CA;MGC142161;MGC142163;PI3K;p110-alpha ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasphosphoinositide-3-kinase, catalytic, alpha polypeptide;PIK3CA;MGC142161;MGC142163;PI3K;p110-alpha ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Uniprot No.	P42336
Species ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasphosphoinositide-3-kinase, catalytic, alpha polypeptide;PIK3CA;MGC142161;MGC142163;PI3K;p110-alpha ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Immunogen	Human PIK3CA
Tested ApplicationsELISA,WBStorage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasphosphoinositide-3-kinase, catalytic, alpha polypeptide;PIK3CA;MGC142161;MGC142163;PI3K;p110-alpha ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Raised In	Rabbit
Storage BufferPBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasphosphoinositide-3-kinase, catalytic, alpha polypeptide;PIK3CA;MGC142161;MGC142163;PI3K;p110-alpha ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Species Reactivity	Human,Mouse,Rat
thaw cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliasphosphoinositide-3-kinase, catalytic, alpha polypeptide;PIK3CA;MGC142161;MGC142163;PI3K;p110-alpha ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Tested Applications	ELISA,WB
IsotypeIgGAliasphosphoinositide-3-kinase, catalytic, alpha polypeptide;PIK3CA;MGC142161;MGC142163;PI3K;p110-alpha ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Storage Buffer	
Aliasphosphoinositide-3-kinase, catalytic, alpha polypeptide;PIK3CA;MGC142161;MGC142163;PI3K;p110-alpha ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Purification Method	Antigen Affinity purified
polypeptide;PIK3CA;MGC142161;MGC142163;PI3K;p110-alpha ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Isotype	IgG
Immunogen SpeciesHomo sapiens (Human)Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Alias	
Target NamesPIK3CATarget DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Product Type	Purified Rabbit Anti human PolyClonal Antibody
Target DetailsPhosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Immunogen Species	Homo sapiens (Human)
a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has	Target Names	PIK3CA
	Target Details	a 110 kDa catalytic subunit. This protein represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has

1