

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

## **PPARG** Antibody

<ul> <li>CSB-PA018424GA01HU</li> <li>Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.</li> <li>P37231</li> <li>Human PPARG</li> <li>Rabbit</li> <li>Human,Mouse,Rat</li> <li>ELISA,WB</li> <li>PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze /</li> </ul>
P37231 Human PPARG Rabbit Human,Mouse,Rat ELISA,WB
Human PPARG Rabbit Human,Mouse,Rat ELISA,WB
Rabbit Human,Mouse,Rat ELISA,WB
Human,Mouse,Rat ELISA,WB
ELISA,WB
PBS with 0.02% Sodium Azide 50% Glycerol pH 7.3 -20°C. Avoid freeze /
thaw cycles.
Antigen Affinity purified
IgG
peroxisome proliferator-activated receptor gamma;PPARG;CIMT1;NR1C3;PPARG1;PPARG2;PPARgamma ;
Purified Rabbit Anti human PolyClonal Antibody
Homo sapiens (Human)
PPARG
This gene encodes a member of the peroxisome proliferator-activated receptor (PPAR) subfamily of nuclear receptors. PPARs form heterodimers with retinoid X receptors (RXRs) and these heterodimers regulate transcription of various genes. Three subtypes of PPARs are known: PPAR-alpha, PPAR-delta, and PPAR-gamma. This protein is PPAR-gamma and is a regulator of adipocyte differentiation. Additionally, PPAR-gamma has been implicated in the pathology of numerous diseases including obesity, diabetes, atherosclerosis and cancer. Alternatively spliced transcript variants that encode different isoforms have been described.

1