



# S100A10 Antibody

|                            |  |
|----------------------------|--|
| <b>Product Code</b>        | CSB-PA020623GA01HU   |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| <b>Uniprot No.</b>         | P60903   |
| <b>Immunogen</b>           | Human S100A10  |
| <b>Raised In</b>           | Rabbit   |
| <b>Species Reactivity</b>  | Human,Mouse,Rat  |
| <b>Tested Applications</b> | ELISA,WB,IHC   |
| <b>Storage Buffer</b>      | PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.3. -20°C, Avoid freeze / thaw cycles.   |
| <b>Purification Method</b> | Antigen Affinity Purified  |
| <b>Isotype</b>             | IgG  |
| <b>Alias</b>               | S100 calcium binding protein<br>A10;S100A10;42C;ANX2L;ANX2LG;CAL1L;CLP11;Ca[1];GP11;MGC111133;P1<br>1;p10 ;  |
| <b>Product Type</b>        | Purified Rabbit Anti human PolyClonal Antibody   |
| <b>Immunogen Species</b>   | Homo sapiens (Human)   |
| <b>Target Names</b>        | S100A10  |
| <b>Target Details</b>      | This protein is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein may function in exocytosis and endocytosis. |
| <b>Usage</b>               | For Research Use Only. Not for use in diagnostic or therapeutic procedures.  |