



# SEC63 Antibody

|                            |  |
|----------------------------|--|
| <b>Product Code</b>        | CSB-PA020961GA01HU   |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| <b>Uniprot No.</b>         | Q9UGP8   |
| <b>Immunogen</b>           | Human SEC63  |
| <b>Raised In</b>           | Rabbit   |
| <b>Species Reactivity</b>  | Human,Mouse,Rat  |
| <b>Tested Applications</b> | ELISA,WB   |
| <b>Storage Buffer</b>      | PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.3. -20°C, Avoid freeze / thaw cycles.   |
| <b>Purification Method</b> | Antigen Affinity purified  |
| <b>Isotype</b>             | IgG  |
| <b>Alias</b>               | SEC63 homolog (S. cerevisiae);SEC63;ERdj2;PRO2507;SEC63L ;   |
| <b>Product Type</b>        | Purified Rabbit Anti human PolyClonal Antibody   |
| <b>Immunogen Species</b>   | Homo sapiens (Human)   |
| <b>Target Names</b>        | SEC63  |
| <b>Target Details</b>      | <p>The Sec61 complex is the central component of the protein translocation apparatus of the endoplasmic reticulum (ER) membrane. This protein and SEC62 protein are found to be associated with ribosome-free SEC61 complex. It is speculated that Sec61-Sec62-Sec63 may perform post-translational protein translocation into the ER. The Sec61-Sec62-Sec63 complex might also perform the backward transport of ER proteins that are subject to the ubiquitin-proteasome-dependent degradation pathway. The encoded protein is an integral membrane protein located in the rough ER.</p> |
| <b>Usage</b>               | For Research Use Only. Not for use in diagnostic or therapeutic procedures.  |