



Recombinant Human 26S proteasome non-ATPase regulatory subunit 4 (PSMD4)

Product Code	CSB-YP018908HU
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P55036
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	>85% (SDS-PAGE)
Sequence	MVLESTMVCV DNSEYMRNGD FLPTRLQAQQ DAVNIVCHSK TRSNPENNVG LITLANDCEV LTTLPDTGR ILSKLHTVQP KGKITFCTGI RVAHLALKHR QGKNHKMR II AFVGGSPVEDN EKDLVKLAKR LKKEKVNVDI INFGEEEVNT EKLTAFVNTL NGKDG TGS HL VTPPPGPSLA DALISSPILA GEGGAMLGLG ASDFEFGVDP SADPELALAL RVSMEEQQR QEEEEARRAAA ASAAEAGIAT TGTEDSDDAL LKMTISQQEF GRTGLPDLSS MTEEEQIAYA MQMSLQGAEF GQAESADIDA SSAMDTSEPA KEEDDYDVMQ DPEFLQSVLE NLPGVDPNNE AIRNAMGSLA SQATKDGKKD KKEEDKK
Source	Yeast
Target Names	PSMD4
Protein Names	Recommended name: 26S proteasome non-ATPase regulatory subunit 4 Alternative name(s): 26S proteasome regulatory subunit RPN10 26S proteasome regulatory subunit S5A Antisecretory factor 1 Short name= AF Short name= ASF Mul
Expression Region	1-377
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full length protein
Target Details	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the non-ATPase subunits of the 19S regulator lid. Pseudogenes have been identified on chromosomes 10 and 21.



Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.