

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com 🍙

WBP1 Antibody

 CSB-PA025974GA01HU Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Q96G27 Human WBP1 Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity Purified IgG
Q96G27 Human WBP1 Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity Purified
 Human WBP1 Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity Purified
Rabbit Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity Purified
 Human,Mouse,Rat ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity Purified
ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity Purified
PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. Antigen Affinity Purified
cycles. Antigen Affinity Purified
5 ,
laG
igo
WW domain binding protein 1;WBP1;MGC15305;WBP-1;
Purified Rabbit Anti human PolyClonal Antibody
Homo sapiens (Human)
WBP1
The globular WW domain is composed of 38 to 40 semiconserved amino acids shared by proteins of diverse functions including structural, regulatory, and signaling proteins. The domain is involved in mediating protein-protein interactions through the binding of polyproline ligands. This gene encodes a WW domain binding protein, which binds to the WW domain of Yes kinase- associated protein by a conserved region: XPPXY motif. The function of this protein has not been determined.

1