




# EIF4A2 Antibody

<b>Product Code</b>	CSB-PA00384A0Rb
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q14240
<b>Immunogen</b>	Recombinant Human Eukaryotic initiation factor 4A-II protein (1-407AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IHC, IF; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200, IF:1:50-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	>95%, Protein G purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Eukaryotic initiation factor 4A-II (eIF-4A-II) (eIF4A-II) (EC 3.6.4.13) (ATP-dependent RNA helicase eIF4A-2), EIF4A2, DDX2B EIF4F
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Epigenetics and Nuclear Signaling
<b>Target Names</b>	EIF4A2

## Image



### Western Blot

Positive WB detected in: HepG2 whole cell lysate, Jurkat whole cell lysate, Raji whole cell lysate, 293 whole cell lysate


All lanes: EIF4A2 antibody at 3µg/ml

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 47 kDa


Observed band size: 47 kDa



Immunohistochemistry of paraffin-embedded human kidney tissue using CSB-PA00384A0Rb at dilution of 1:100



Immunohistochemistry of paraffin-embedded human breast cancer using CSB-PA00384A0Rb at dilution of 1:100



Immunofluorescence staining of A549 cells with CSB-PA00384A0Rb at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

**Usage**

For Research Use Only. Not for use in diagnostic or therapeutic procedures.