



Recombinant Human Mitogen-activated protein kinase 10 (MAPK10)

Product Code	CSB-BP013449HU
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P53779
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	>85% (SDS-PAGE)
Sequence	MSLHFLYYCS EPTLDVKIAF CQGFDKQVDV SYIAKHYNMS KSKVDNQFYS VEVGDSTFTV LKRYQNLKPI GSGAQQIVCA AYDAVLDRNV AIKKLSRPFQ NQTHAKRAYR ELVLMKCVNH KNIISLLNVF TPQKTLEEFQ DVYLVME LMD ANLCQVIQME LDHERMSYLL YQMLCGIKHL HSAGIIHRDL KPSNIVVKSD CTLKILDFGL ARTAGTSFMM TPYVVTRYR APEVILGMGY KENVDIWSVG CIMGEMVRHK ILFPGRDYID QWNKVIEQLG TPCPEFMKKL QPTVRNYVEN RPKYAGLTFP KLFPDSLFP DSEHNKLGAS QARDLLSKML VIDPAKRISV DDALQHPYIN VWYDPAEVEA PPPQIYDKQL DEREHTIEEW KELIYKEVMN SEEKTKNGVV KGQPSPSGAA VNSSESLPPS SSVNDISSMS TDQTLASD TD SSLEASAGPL GCCR
Source	Baculovirus
Target Names	MAPK10
Protein Names	Recommended name: Mitogen-activated protein kinase 10 Short name= MAP kinase 10 Short name= MAPK 10 EC= 2.7.11.24 Alternative name(s): MAP kinase p49 3F12 Stress-activated protein kinase 1b Short name= SAPK1b St
Expression Region	1-464
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full length protein
Target Details	This protein is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This protein is a neuronal-specific form of c-Jun N-terminal kinases (JNKs). Through its phosphorylation and nuclear localization, this kinase plays regulatory roles in the signaling pathways during neuronal apoptosis. Beta-arrestin 2, a receptor-regulated MAP kinase scaffold protein, is found to interact with, and stimulate the phosphorylation of this kinase by MAP kinase kinase 4 (MKK4). Cyclin-dependent kinase 5 can phosphorylate, and inhibit the activity of this kinase, which may be important in preventing neuronal apoptosis. Four alternatively spliced transcript variants encoding distinct



isoforms have been reported.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.